

# **DBH Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20809c

#### **Product Information**

**Application** WB, E **Primary Accession** P09172 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB50735 Calculated MW 69065

#### **Additional Information**

Gene ID 1621

Other Names Dopamine beta-hydroxylase, Dopamine beta-monooxygenase, Soluble

dopamine beta-hydroxylase, DBH

**Target/Specificity**This DBH antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 199-232 amino acids from the Central

region of human DBH.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** DBH Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name DBH

**Function** Catalyzes the hydroxylation of dopamine to noradrenaline (also known as

norepinephrine), and is thus vital for regulation of these neurotransmitters.

**Cellular Location** [Soluble dopamine beta-hydroxylase]: Cytoplasmic vesicle, secretory vesicle

lumen Cytoplasmic vesicle, secretory vesicle, chromaffin granule lumen.


## **Background**

Conversion of dopamine to noradrenaline.

### References

Humphray S.J., et al. Nature 429:369-374(2004). Lamouroux A., et al. EMBO J. 6:3931-3937(1987). Kobayashi K., et al. Nucleic Acids Res. 17:1089-1102(1989). Li B., et al. Biochem. J. 313:57-64(1996). Liu T., et al. J. Proteome Res. 4:2070-2080(2005).

## **Images**



Western blot analysis of lysate from SH-SY5Y cell line, using DBH Antibody (Center)(Cat. #AP20809c). AP20809c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.