

# QTRT1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20812c

## Product Information

Application	WB, E
Primary Accession	<a href="#">Q9BXR0</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50086
Calculated MW	44048

## Additional Information

Gene ID	81890
Other Names	Queueine tRNA-ribosyltransferase, Guanine insertion enzyme, tRNA-guanine transglycosylase, QTRT1, TGT, TGUT
Target/Specificity	This QTRT1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 303-337 amino acids from the C-terminal region of human QTRT1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	QTRT1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	QTRT1 {ECO:0000255 HAMAP-Rule:MF_03218}
Synonyms	TGT, TGUT
Function	Catalytic subunit of the queueine tRNA-ribosyltransferase (TGT) that catalyzes the base-exchange of a guanine (G) residue with queueine (Q) at position 34 (anticodon wobble position) in tRNAs with GU(N) anticodons (tRNA-Asp, -Asn,

-His and -Tyr), resulting in the hypermodified nucleoside queuosine (7-(((4,5-cis-dihydroxy-2-cyclopenten-1-yl)amino)methyl)-7-deazaguanosine) (PubMed:[11255023](#), PubMed:[20354154](#), PubMed:[34009357](#), PubMed:[34241577](#)). Catalysis occurs through a double-displacement mechanism. The nucleophile active site attacks the C1' of nucleotide 34 to detach the guanine base from the RNA, forming a covalent enzyme-RNA intermediate. The proton acceptor active site deprotonates the incoming queuine, allowing a nucleophilic attack on the C1' of the ribose to form the product (By similarity). Modification of cytoplasmic tRNAs with queuosine controls the elongation speed of cognate codons, thereby ensuring the correct folding of nascent proteins to maintain proteome integrity (PubMed:[30093495](#)).

### Cellular Location

Cytoplasm {ECO:0000255|HAMAP-Rule:MF\_03218}. Mitochondrion outer membrane {ECO:0000255|HAMAP-Rule:MF\_03218}; Peripheral membrane protein {ECO:0000255|HAMAP-Rule:MF\_03218}; Cytoplasmic side {ECO:0000255|HAMAP-Rule:MF\_03218}. Note=Weakly associates with mitochondria, possibly via QTRT2. {ECO:0000255|HAMAP-Rule:MF\_03218}


## Background

Interacts with QTRTD1 to form an active queuine tRNA- ribosyltransferase. This enzyme exchanges queuine for the guanine at the wobble position of tRNAs with GU(N) anticodons (tRNA-Asp, -Asn, -His and -Tyr), thereby forming the hypermodified nucleoside queuosine (Q) (7-(((4,5-cis-dihydroxy-2-cyclopenten-1-yl)amino)methyl)-7-deazaguanosine) (By similarity).

## References

Deshpande K.L.,et al.Gene 265:205-212(2001).  
Grimwood J.,et al.Nature 428:529-535(2004).  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).  
Van Damme P.,et al.Proc. Natl. Acad. Sci. U.S.A. 109:12449-12454(2012).

## Images



Western blot analysis of lysate from human heart tissue lysate, using QTRT1 Antibody (C-term)(Cat. #AP20812c). AP20812c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.