

# Mouse Cbx7 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20840c

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q8VDS3</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB50982
<b>Calculated MW</b>	18109

## Additional Information

---

<b>Gene ID</b>	52609
<b>Other Names</b>	Chromobox protein homolog 7, Cbx7, D15Ert417e
<b>Target/Specificity</b>	This Mouse Cbx7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 144-178 amino acids from the C-terminal region of human Mouse Cbx7.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Mouse Cbx7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	Cbx7
<b>Synonyms</b>	D15Ert417e
<b>Function</b>	Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed: <a href="#">16537902</a> , PubMed: <a href="#">22226355</a> ). PcG PRC1 complex acts via

chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Promotes histone H3 trimethylation at 'Lys-9' (H3K9me3) (By similarity). Binds to histone H3 trimethylated at 'Lys-9' (H3K9me3) or at 'Lys-27' (H3K27me3) (PubMed:[16537902](#), PubMed:[22226355](#)). Trimethylation at 'Lys-27' (H3K27me3) is important for chromatin recruitment (PubMed:[16537902](#), PubMed:[22226355](#)). May possibly also bind trimethylated lysine residues in other proteins (in vitro) (PubMed:[16537902](#)). Binds non-coding, single-stranded RNA and double-stranded RNA (PubMed:[16537902](#), PubMed:[20541999](#)). Plays a role in the timely repression of differentiation-specific genes in pluripotent embryonic stem cells to maintain the undifferentiated state (PubMed:[22226355](#)). Regulator of cellular lifespan by maintaining the repression of CDKN2A, but not by inducing telomerase activity (PubMed:[14647293](#)).

## Cellular Location

Nucleus. Chromosome Note=Requires trimethylation at 'Lys-27' (H3K27me3) for the localization to chromatin (PubMed:[22226355](#)). Localizes to facultative heterochromatin and to the inactivated X chromosome in females (PubMed:[16537902](#)).

## Tissue Location

Expressed in embryonic stem cells.


## Background

Component of a Polycomb group (PcG) multiprotein PRC1- like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Promotes histone H3 trimethylation at 'Lys-9' (H3K9me3). Binds to histone H3 trimethylated 'Lys-9' (H3K9me3) or at 'Lys-27' (H3K27me3). May possibly also bind trimethylated lysine residues in other proteins (in vitro). Binds non-coding, single-stranded RNA. Regulator of cellular lifespan by maintaining the repression of CDKN2A, but not by inducing telomerase activity.

## References

Gil J.,et al.Nat. Cell Biol. 6:67-72(2004).  
Bernstein E.,et al.Mol. Cell. Biol. 26:2560-2569(2006).  
Yap K.L.,et al.Mol. Cell 38:662-674(2010).

## Images



Western blot analysis of lysates from HeLa cell line, mouse brain, mouse testis, mouse lung, mouse spleen, rat brain tissue (from left to right), using Mouse Cbx7 Antibody (C-term)(Cat. #AP20840c). AP20840c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.