

SERPINE1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20852c

Product Information

Application	WB, E
Primary Accession	P05121
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB30793
Calculated MW	45060

Additional Information

Gene ID	5054
Other Names	Plasminogen activator inhibitor 1, PAI, PAI-1, Endothelial plasminogen activator inhibitor, Serpin E1, SERPINE1, PAI1, PLANH1
Target/Specificity	This SERPINE1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 323-357 amino acids from the C-terminal region of human SERPINE1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SERPINE1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SERPINE1
Synonyms	PAI1, PLANH1
Function	Serine protease inhibitor. Inhibits TMPRSS7 (PubMed: 15853774). Is a primary inhibitor of tissue-type plasminogen activator (PLAT) and urokinase-type plasminogen activator (PLAU). As PLAT inhibitor, it is required

for fibrinolysis down-regulation and is responsible for the controlled degradation of blood clots (PubMed:[17912461](#), PubMed:[8481516](#), PubMed:[9207454](#), PubMed:[21925150](#)). As PLAU inhibitor, it is involved in the regulation of cell adhesion and spreading (PubMed:[9175705](#)). Acts as a regulator of cell migration, independently of its role as protease inhibitor (PubMed:[15001579](#), PubMed:[9168821](#)). It is required for stimulation of keratinocyte migration during cutaneous injury repair (PubMed:[18386027](#)). It is involved in cellular and replicative senescence (PubMed:[16862142](#)). Plays a role in alveolar type 2 cells senescence in the lung (By similarity). Is involved in the regulation of cementogenic differentiation of periodontal ligament stem cells, and regulates odontoblast differentiation and dentin formation during odontogenesis (PubMed:[25808697](#), PubMed:[27046084](#)).

Cellular Location

Secreted.

Tissue Location

Expressed in endothelial cells (PubMed:2430793, PubMed:3097076). Found in plasma, platelets, and hepatoma and fibrosarcoma cells.


Background

Serine protease inhibitor. This inhibitor acts as 'bait' for tissue plasminogen activator, urokinase, protein C and matriptase-3/TMPRSS7. Its rapid interaction with PLAT may function as a major control point in the regulation of fibrinolysis.

References

Pannekoek H.,et al.EMBO J. 5:2539-2544(1986).
Loskutoff D.J.,et al.Biochemistry 26:3763-3768(1987).
Ginsburg D.,et al.J. Clin. Invest. 78:1673-1680(1986).
Follo M.,et al.Gene 84:447-453(1989).
Strandberg L.,et al.Eur. J. Biochem. 176:609-616(1988).

Images



Western blot analysis of lysate from human liver tissue, using SERPINE1 Antibody (C-term)(Cat. #AP20852c). AP20852c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.