

# CTDSP1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20906a

# **Product Information**

Application	WB, E
Primary Accession	<u>Q9GZU7</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50787
Calculated MW	29203

# **Additional Information**

Gene ID	58190
Other Names	Carboxy-terminal domain RNA polymerase II polypeptide A small phosphatase 1, Nuclear LIM interactor-interacting factor 3, NLI-IF, NLI-interacting factor 3, Small C-terminal domain phosphatase 1, SCP1, Small CTD phosphatase 1, CTDSP1, NIF3, NLIIF, SCP1
Target/Specificity	This CTDSP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 96-130 amino acids from the Central region of human CTDSP1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTDSP1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name	CTDSP1
Synonyms	NIF3, NLIIF, SCP1
Function	Preferentially catalyzes the dephosphorylation of 'Ser-5' within the tandem

	7 residue repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.
Cellular Location	Nucleus. Note=Colocalizes with RNA polymerase II
Tissue Location	Expression is restricted to non-neuronal tissues. Highest expression in skeletal muscle, spleen, lung and placenta

## Background

Preferentially catalyzes the dephosphorylation of 'Ser- 5' within the tandem 7 residues repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.

## References

Marquet S.,et al.Mamm. Genome 11:755-762(2000). Yeo M.,et al.J. Biol. Chem. 278:26078-26085(2003). Li W.B.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases. Hillier L.W.,et al.Nature 434:724-731(2005). Yeo M.,et al.Science 307:596-600(2005).

### Images



Western blot analysis of lysates from HepG2, Jurkat cell line, mouse liver tissue lysate (from left to right), using CTDSP1 Antibody (Center)(Cat. #AP20906a). AP20906a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.