10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CTDSP1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20906a

Product Information

Application	WB, E
Primary Accession	Q9GZU7
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50787
Calculated MW	29203

Additional Information

Gene ID	58190
Other Names	Carboxy-terminal domain RNA polymerase II polypeptide A small phosphatase 1, Nuclear LIM interactor-interacting factor 3, NLI-IF, NLI-interacting factor 3, Small C-terminal domain phosphatase 1, SCP1, Small CTD phosphatase 1, CTDSP1, NIF3, NLIIF, SCP1
Target/Specificity	This CTDSP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 96-130 amino acids from the Central region of human CTDSP1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTDSP1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CTDSP1
Synonyms	NIF3, NLIIF, SCP1
Function	Preferentially catalyzes the dephosphorylation of 'Ser-5' within the tandem

	7 residue repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.
Cellular Location	Nucleus. Note=Colocalizes with RNA polymerase II
Tissue Location	Expression is restricted to non-neuronal tissues. Highest expression in skeletal muscle, spleen, lung and placenta


Background

Preferentially catalyzes the dephosphorylation of 'Ser- 5' within the tandem 7 residues repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.

References

Marquet S.,et al.Mamm. Genome 11:755-762(2000).
Yeo M.,et al.J. Biol. Chem. 278:26078-26085(2003).
Li W.B.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.
Hillier L.W.,et al.Nature 434:724-731(2005).
Yeo M.,et al.Science 307:596-600(2005).

Images



Western blot analysis of lysates from HepG2, Jurkat cell line, mouse liver tissue lysate (from left to right), using CTDSP1 Antibody (Center)(Cat. #AP20906a). AP20906a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.