

Mouse Dbx1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20920c

Product Information

Application WB, E Primary Accession P52950

Reactivity Human, Rat, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB51057
Calculated MW 36334

Additional Information

Gene ID 13172

Other Names Homeobox protein DBX1, Developing brain homeobox protein 1, Dbx1, Dbx

Target/Specificity This Mouse Dbx1 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 261-295 amino acids from the

C-terminal region of Mouse Dbx1.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Mouse Dbx1 Antibody (C-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name Dbx1

Synonyms Dbx

Function Could have a role in patterning the central nervous system during

embryogenesis. Has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons.

Regulates the transcription factor profile, neurotransmitter phenotype,

intraspinal migratory path and axonal trajectory of V0 neurons, features that differentiate them from an adjacent set of V1 neurons.

Cellular Location

Nucleus.


Background

Could have a role in patterning the central nervous system during embryogenesis. Has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons. Regulates the transcription factor profile, neurotransmitter phenotype, intraspinal migratory path and axonal trajectory of V0 neurons, features that differentiate them from an adjacent set of V1 neurons.

References

Lu S., et al. Mech. Dev. 47:187-195(1994). Carninci P., et al. Science 309:1559-1563(2005). Pierani A., et al. Neuron 29:367-384(2001).

Images



Western blot analysis of lysate from mouse skeletal muscle tissue, using Dbx1 Antibody (C-term)(Cat. #AP20920c). AP20920c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.