

# Mouse Ddr2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20936a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q62371</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB50401

## Additional Information

---

<b>Other Names</b>	Discoidin domain-containing receptor 2, Discoidin domain receptor 2, CD167 antigen-like family member B, Neurotrophic tyrosine kinase, receptor-related 3, Receptor protein-tyrosine kinase TKT, Tyrosine-protein kinase TYRO10, CD167b, Ddr2, Ntrkr3, Tkt, Tyro10
<b>Target/Specificity</b>	This Mouse Ddr2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 503-537 amino acids from the Central region of Mouse Ddr2.
<b>Dilution</b>	WB--1:1000 E--Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Mouse Ddr2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

### Background

---

Tyrosine kinase that functions as cell surface receptor for fibrillar collagen and regulates cell differentiation, remodeling of the extracellular matrix, cell migration and cell proliferation. Required for normal bone development. Regulates osteoblast differentiation and chondrocyte maturation via a signaling pathway that involves MAP kinases and leads to the activation of the transcription factor RUNX2. Regulates remodeling of the extracellular matrix by up-regulation of the collagenases MMP1, MMP2 and MMP13, and

thereby facilitates cell migration and tumor cell invasion. Promotes fibroblast migration and proliferation, and thereby contributes to cutaneous wound healing.

## References

---

Karn T.,et al.Oncogene 8:3433-3440(1993).

Lai C.,et al.Oncogene 9:877-883(1994).


Labrador J.P.,et al.EMBO Rep. 2:446-452(2001).

Olaso E.,et al.J. Biol. Chem. 277:3606-3613(2002).

Ikeda K.,et al.J. Biol. Chem. 277:19206-19212(2002).

## Images

---



Western blot analysis of lysates from mouse NIH/3T3 cell line and mouse lung tissue(from left to right), using Ddr2 Antibody (Center)(Cat. #AP20936a). AP20936a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.