

# EWSR1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20981a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q01844</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB51592
<b>Calculated MW</b>	68478

## Additional Information

---

<b>Gene ID</b>	2130
<b>Other Names</b>	RNA-binding protein EWS, EWS oncogene, Ewing sarcoma breakpoint region 1 protein, EWSR1, EWS
<b>Target/Specificity</b>	This EWSR1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 328-362 amino acids from the Central region of human EWSR1.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	EWSR1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	EWSR1
<b>Synonyms</b>	EWS
<b>Function</b>	Binds to ssRNA containing the consensus sequence 5'-AGGUAA-3' (PubMed: <a href="#">21256132</a> ). Might normally function as a transcriptional repressor (PubMed: <a href="#">10767297</a> ). EWS-fusion-proteins (EFPS) may play a role in the

tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

**Cellular Location** Nucleus. Cytoplasm. Cell membrane. Note=Relocates from cytoplasm to ribosomes upon PTK2B/FAK2 activation

**Tissue Location** Ubiquitous.

## Background

---

Might normally function as a transcriptional repressor. EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.


## References

---

Delattre O., et al. Nature 359:162-165(1992).  
Plougastel B., et al. Genomics 18:609-615(1993).  
Zucman-Rossi J., et al. Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.  
Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004).  
Ota T., et al. Nat. Genet. 36:40-45(2004).

## Images

---



Western blot analysis of lysate from mouse NIH/3T3 cell line, using EWSR1 Antibody (Center)(Cat. #AP20981a). AP20981a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.