

NFIA Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20998c

Product Information

Application	WB, E
Primary Accession	Q12857
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB51284
Calculated MW	55944

Additional Information

Gene ID	4774
Other Names	Nuclear factor 1 A-type, NF1-A, Nuclear factor 1/A, CCAAT-box-binding transcription factor, CTF, Nuclear factor I/A, NF-I/A, NFI-A, TGGCA-binding protein, NFIA, KIAA1439
Target/Specificity	This NFIA antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 384-418 amino acids from the C-terminal region of human NFIA.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NFIA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NFIA
Synonyms	KIAA1439
Function	Recognizes and binds the palindromic sequence 5'- TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of

adenovirus type 2. These proteins are individually capable of activating transcription and replication.

Cellular Location Nucleus.


Background

Recognizes and binds the palindromic sequence 5'- TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

References

Nagase T.,et al.DNA Res. 7:65-73(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Qian F.,et al.Genomics 28:66-73(1995).

Images



Western blot analysis of lysates from Hela cell line, rat liver tissue lysate(from left to right), using NFIA Antibody (C-term)(Cat. #AP20998c). AP20998c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Citations

- [MicroRNA 373 Facilitates the Replication of Porcine Reproductive and Respiratory Syndrome Virus by Its Negative Regulation of Type I Interferon Induction.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.