

NACC1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21031a

Product Information

Application	WB, E
Primary Accession	Q96RE7
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB51582

Additional Information

Other Names	Nucleus accumbens-associated protein 1, NAC-1, BTB/POZ domain-containing protein 14B, NACC1, BTBD14B, NAC1
Target/Specificity	This NACC1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 324-358 amino acids from the Central region of human NACC1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NACC1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Background

Functions as a transcriptional repressor. Seems to function as a transcriptional corepressor in neuronal cells through recruitment of HDAC3 and HDAC4. Contributes to tumor progression, and tumor cell proliferation and survival. This may be mediated at least in part through repressing transcriptional activity of GADD45GIP1. Required for recruiting the proteasome from the nucleus to the cytoplasm and dendritic spines.


References

Cha X.Y.,et al.Submitted (JUN-2001) to the EMBL/GenBank/DDBJ databases.
Nakayama K.,et al.Proc. Natl. Acad. Sci. U.S.A. 103:18739-18744(2006).
Nakayama K.,et al.Cancer Res. 67:8058-8064(2007).
Davidson B.,et al.Hum. Pathol. 38:1030-1036(2007).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

Images



Anti-NACC1 Antibody (Center)at 1:1000 dilution + human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of lysate from mouse kidney tissue, using NACC1 Antibody (Center)(Cat. #AP21031a). AP21031a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.