10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# (Mouse) Epcam Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21054a

## Product Information

| | |
|---|---|
| **Application** | WB, FC, E |
| **Primary Accession** | Q99JW5 |
| **Reactivity** | Human, Rat, Mouse |
| **Predicted** | Bovine, Canine |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB51264 |
| **Calculated MW** | 35019 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 17075 |
| **Other Names** | Epithelial cell adhesion molecule, Ep-CAM, Epithelial glycoprotein 314, EGP314, mEGP314, Protein 289A, Tumor-associated calcium signal transducer 1, CD326, Epcam, Tacstd1 |
| **Target/Specificity** | This Mouse Epcam antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 302-335 amino acids from the C-terminal region of mouse Epcam. |
| **Dilution** | WB~~1:1000 FC~~1:25 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | (Mouse) Epcam Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | Epcam |
| **Synonyms** | Tacstd1 |
| **Function** | May act as a physical homophilic interaction molecule between intestinal |

| | |
|---|---|
| | epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E (By similarity). |
| **Cellular Location** | Lateral cell membrane {ECO:0000250\|UniProtKB:P16422}; Single-pass type I membrane protein {ECO:0000250\|UniProtKB:P16422}. Cell junction, tight junction {ECO:0000250\|UniProtKB:P16422}. Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction {ECO:0000250\|UniProtKB:P16422} |


## Background

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E (By similarity).


## References

Bergsagel P.L.,et al.J. Immunol. 148:590-596(1992).
Carninci P.,et al.Science 309:1559-1563(2005).

## Images




Flow cytometric analysis of HepG2 cells using (Mouse) Epcam Antibody (C-term)(green, Cat#AP21054a) compared to an isotype control of rabbit IgG(blue). AP21054a was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit lgG at 1:400 dilution was used as the secondary antibody.




Western blot analysis of lysates from mouse kidney, mouse lung, mouse testis tissue lysate (from left to right), using Epcam Antibody (C-term)(Cat. #AP21054a). AP21054a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

## Citations

- Extracellular Vesicles Enriched with Moonlighting Metalloproteinase Are Highly Transmissive, Pro-Tumorigenic, and Trans-Activates Cellular Communication Network Factor ( ): CRISPR against Cancer

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.