

HEMGN Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21062a

Product Information

Application	WB, E
Primary Accession	Q9BXL5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB51964
Calculated MW	55341

Additional Information

Gene ID	55363
Other Names	Hemogen, Erythroid differentiation-associated gene protein, EDAG-1, Hemopoietic gene protein, Negative differentiation regulator protein, HEMGN, EDAG, NDR
Target/Specificity	This HEMGN antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 355-389 amino acids from the C-terminal region of human HEMGN.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HEMGN Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HEMGN
Synonyms	EDAG, NDR
Function	Regulates the proliferation and differentiation of hematopoietic cells. Overexpression block the TPA-induced megakaryocytic differentiation in the

K562 cell model. May also prevent cell apoptosis through the activation of the nuclear factor-kappa B (NF-kB).

Cellular Location

Nucleus.

Tissue Location

Expressed in hematopoietic precursor cells, thyroid and spermatids (at protein level). Expressed in bone marrow, testis, thymus. Expressed in prostate cancer and ovarian cancer. Also expressed in thymus and thyroid tumors, non-Hodgkin lymphoma, various leukemia cell lines, peripheral blood mononuclear cells (PBMCs) and bone marrow mononuclear cells (BMMCs) of patients with leukemia


Background

Regulates the proliferation and differentiation of hematopoietic cells. Overexpression block the TPA-induced megakaryocytic differentiation in the K562 cell model. May also prevent cell apoptosis through the activation of the nuclear factor-kappa B (NF-kB).

References

Yang L.V.,et al.Mech. Dev. 104:105-111(2001).
Yang L.V.,et al.Dev. Dyn. 228:606-616(2003).
Li C.Y.,et al.Cell Death Differ. 11:1299-1308(2004).
Liu C.C.,et al.J. Biomed. Sci. 11:104-116(2004).
Humphray S.J.,et al.Nature 429:369-374(2004).

Images



Anti-HEMGN Antibody (C-term) at 1:2000 dilution + Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.