

MAOA Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21078a

Product Information

Application	WB, E
Primary Accession	P21397
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB48811
Calculated MW	59682

Additional Information

Gene ID	4128
Other Names	Amine oxidase [flavin-containing] A, Monoamine oxidase type A, MAO-A, MAOA
Target/Specificity	This MAOA antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 465-499 amino acids from the C-terminal region of human MAOA.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MAOA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MAOA (HGNC:6833)
Function	Catalyzes the oxidative deamination of primary and some secondary amine such as neurotransmitters, with concomitant reduction of oxygen to hydrogen peroxide and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed: 18391214 , PubMed: 20493079 , PubMed: 24169519 ,

PubMed:[8316221](#)). Preferentially oxidizes serotonin (PubMed:[20493079](#), PubMed:[24169519](#)). Also catalyzes the oxidative deamination of kynuramine to 3-(2-aminophenyl)-3-oxopropanal that can spontaneously condense to 4-hydroxyquinoline (By similarity).

Cellular Location	Mitochondrion outer membrane {ECO:0000250 UniProtKB:P21396}; Single-pass type IV membrane protein {ECO:0000250 UniProtKB:P21396}; Cytoplasmic side {ECO:0000250 UniProtKB:P21396}
Tissue Location	Heart, liver, duodenum, blood vessels and kidney.


Background

Catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

References

Hsu Y.-P.P.,et al.J. Neurochem. 51:1321-1324(1988).
Bach A.W.J.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:4934-4938(1988).
Chen Z.-Y.,et al.Nucleic Acids Res. 19:4537-4541(1991).
Grimsby J.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:3637-3641(1991).
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of lysate from human placenta tissue lysate, using MAOA Antibody (C-term)(Cat. #AP21078a). AP21078a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.