

CSK Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21135a

Product Information

Application	WB, E
Primary Accession	<u>P41240</u>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB50499
Calculated MW	50704

Additional Information

Gene ID	1445
Other Names	Tyrosine-protein kinase CSK, C-Src kinase, Protein-tyrosine kinase CYL, CSK
Target/Specificity	This CSK antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 164-197 amino acids from the Central region of human CSK.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CSK Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name CSK
Function Non-receptor tyrosine-protein kinase that plays an importan regulation of cell growth, differentiation, migration and immu Phosphorylates tyrosine residues located in the C- terminal ta kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1 phosphorylation, Src-family members engage in intramolecul between the phosphotyrosine tail and the SH2 domain that re

	inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.
Cellular Location	Cytoplasm. Cell membrane. Note=Mainly cytoplasmic, also present in lipid rafts
Tissue Location	Expressed in lung and macrophages.

Background

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T- cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.

References

Partanen J., et al.Oncogene 6:2013-2018(1991). Braeuninger A., et al.Proc. Natl. Acad. Sci. U.S.A. 88:10411-10415(1991). Brauninger A., et al.Gene 110:205-211(1992). Braeuninger A., et al.Oncogene 8:1365-1369(1993). Halleck A., et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of lysate from Ramos cell line, using CSK Antibody (Center)(Cat. #AP21135a). AP21135a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.