

FGF9 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21143a

Product Information

Application	WB, IHC, FC, E
Primary Accession	P31371
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB51887
Calculated MW	23441

Additional Information

Gene ID	2254
Other Names	Fibroblast growth factor 9, FGF-9, Glia-activating factor, GAF, Heparin-binding growth factor 9, HBGF-9, FGF9
Target/Specificity	This FGF9 antibody is generated from a rabbit immunized with a recombinant protein of human FGF9.
Dilution	WB~~1:4000 IHC~~1:100~500 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FGF9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FGF9
Function	Plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. May have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.

Cellular Location	Secreted.
Tissue Location	Glial cells.


Background

Plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. May have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.


References

Miyamoto M.,et al.Mol. Cell. Biol. 13:4251-4259(1993).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham A.,et al.Nature 428:522-528(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Naruo K.,et al.J. Biol. Chem. 268:2857-2864(1993).

Images





Overlay histogram showing MCF-7 cells stained with AP21143a (green line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (1583138) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



AP21143a staining FGF9 in Human liver tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

All lanes : Anti-FGF9 Antibody at 1:4000 dilution Lane 1: SW480 whole cell lysates Lane 2: mouse kidney lysates Lane 3: human brain lysates Lane 4: human kidney lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of lysate from human kidney tissue lysate, using FGF9 Antibody (Cat. #AP21143a). AP21143a was diluted at 1:2000. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.