10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NRBP2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21153a

Product Information

Application	WB, E
Primary Accession	Q9NSY0
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB50918
Calculated MW	57803

Additional Information

Gene ID	340371
Other Names	Nuclear receptor-binding protein 2, Transformation-related gene 16 protein, TRG-16, NRBP2 (HGNC:19339)
Target/Specificity	This NRBP2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 394-428 amino acids from the C-terminal region of human NRBP2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NRBP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NRBP2 (HGNC:19339)
Function	May regulate apoptosis of neural progenitor cells during their differentiation.
Cellular Location	Cytoplasm.


Background

May regulate apoptosis of neural progenitor cells during their differentiation.

References

Kim J.W.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Wan D.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:15724-15729(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Nusbaum C.,et al.Nature 439:331-335(2006).

Images



All lanes: Anti-NRBP2 Antibody (C-term) at 1:500 dilution Lane 1: U-251 MG whole cell lysate Lane 2: Mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 60, 35 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.