

SSTR1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21173a

Product Information

Application	WB, E
Primary Accession	P30872
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52188
Calculated MW	42686

Additional Information

Gene ID	6751
Other Names	Somatostatin receptor type 1, SS-1-R, SS1-R, SS1R, SRIF-2, SSTR1
Target/Specificity	This SSTR1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 367-403 amino acids from the C-terminal region of human SSTR1.
Dilution	WB~~1:4000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SSTR1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SSTR1
Function	Receptor for somatostatin with higher affinity for somatostatin-14 than -28. This receptor is coupled via pertussis toxin sensitive G proteins to inhibition of adenylyl cyclase. In addition it stimulates phosphotyrosine phosphatase and Na(+)/H(+) exchanger via pertussis toxin insensitive G proteins.
Cellular Location	Cell membrane; Multi-pass membrane protein.

Tissue Location

Fetal kidney, fetal liver, and adult pancreas, brain, lung, jejunum and stomach

Background

Receptor for somatostatin with higher affinity for somatostatin-14 than -28. This receptor is coupled via pertussis toxin sensitive G proteins to inhibition of adenylyl cyclase. In addition it stimulates phosphotyrosine phosphatase and Na(+)/H(+) exchanger via pertussis toxin insensitive G proteins.


References

Yamada Y.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:251-255(1992).


Kopatz S.A.,et al.Submitted (JUN-2003) to the EMBL/GenBank/DDBJ databases.

Schwaerzler A.,et al.J. Biol. Chem. 275:9557-9562(2000).

Images



Anti-SSTR1 Antibody (C-term) at 1:4000 dilution + THP-1 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-SSTR1 Antibody (C-term) at 1:4000 dilution
Lane 1: human pancreas lysates Lane 2: human lung lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.