

# UBE2L3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2117b

## Product Information

---

<b>Application</b>	IHC-P, WB, E
<b>Primary Accession</b>	<a href="#">P68036</a>
<b>Other Accession</b>	<a href="#">P68037</a> , <a href="#">Q3MHP1</a> , <a href="#">NP_003338</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Predicted</b>	Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	17862
<b>Antigen Region</b>	123-153

## Additional Information

---

<b>Gene ID</b>	7332
<b>Other Names</b>	Ubiquitin-conjugating enzyme E2 L3, L-UBC, Ubch7, Ubiquitin carrier protein L3, Ubiquitin-conjugating enzyme E2-F1, Ubiquitin-protein ligase L3, UBE2L3, UBCE7, UBCH7
<b>Target/Specificity</b>	This UBE2L3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 123-153 amino acids from the C-terminal region of human UBE2L3.
<b>Dilution</b>	IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	UBE2L3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	UBE2L3
<b>Synonyms</b>	UBCE7, UBCH7

<b>Function</b>	Ubiquitin-conjugating enzyme E2 that specifically acts with HECT-type and RBR family E3 ubiquitin-protein ligases. Does not function with most RING-containing E3 ubiquitin-protein ligases because it lacks intrinsic E3-independent reactivity with lysine; in contrast, it has activity with the RBR family E3 enzymes, such as PRKN, RNF31 and ARIH1, that function like RING-HECT hybrids. Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. Mediates ubiquitination by the CUL9-RBX1 complex (PubMed: <a href="#">38605244</a> ). In vitro catalyzes 'Lys-11'-linked polyubiquitination. Involved in the selective degradation of short-lived and abnormal proteins. Down-regulated during the S-phase it is involved in progression through the cell cycle. Regulates nuclear hormone receptors transcriptional activity. May play a role in myelopoiesis.
<b>Cellular Location</b>	Nucleus. Cytoplasm
<b>Tissue Location</b>	Ubiquitous, with highest expression in testis.


## Background

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). UBE2L3 is a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is demonstrated to participate in the ubiquitination of p53, c-Fos, and the NF- $\kappa$ B precursor p105 in vitro.

## References


Moynihan, T.P., et al., Genomics 51(1):124-127 (1998).  
 Moynihan, T.P., et al., Mamm. Genome 7(7):520-525 (1996).  
 Nuber, U., et al., J. Biol. Chem. 271(5):2795-2800 (1996).  
 Robinson, P.A., et al., Mamm. Genome 6(10):725-731 (1995).  
 Ardley, H.C., et al., Biochim. Biophys. Acta 1491 (1-3), 57-64 (2000).

## Images



UBE2L3 Antibody (C-term)(Cat. #AP2117b) western blot analysis in U87-MG,mouse NIH/3T3, rat PC-12 cell line and mouse testis, rat testis tissue lysates (35ug/lane). This demonstrates the UBE2L3 antibody detected the UBE2L3 protein (arrow).

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.