

# CNR2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21180a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P34972</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB52190
<b>Calculated MW</b>	39681

## Additional Information

---

<b>Gene ID</b>	1269
<b>Other Names</b>	Cannabinoid receptor 2, CB-2, CB2, hCB2, CX5, CNR2, CB2A, CB2B
<b>Target/Specificity</b>	This CNR2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 18-53 amino acids from the N-terminal region of human CNR2.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CNR2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CNR2
<b>Synonyms</b>	CB2A, CB2B
<b>Function</b>	Heterotrimeric G protein-coupled receptor for endocannabinoid 2-arachidonoylglycerol mediating inhibition of adenylate cyclase. May function in inflammatory response, nociceptive transmission and bone homeostasis.

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite. Perikaryon Note=Localizes to apical dendrite of pyramidal neurons.

**Tissue Location**


Preferentially expressed in cells of the immune system with higher expression in B-cells and NK cells (at protein level). Expressed in skin in suprabasal layers and hair follicles (at protein level). Highly expressed in tonsil and to a lower extent in spleen, peripheral blood mononuclear cells, and thymus. PubMed:14657172 could not detect expression in normal brain. Expressed in brain by perivascular microglial cells and dorsal root ganglion sensory neurons (at protein level). Two isoforms are produced by alternative promoter usage and differ only in the 5' UTR: isoform CB2A is observed predominantly in testis with some expression in brain, while isoform CB2B is predominant in spleen and leukocytes

**Background**

Heterotrimeric G protein-coupled receptor for endocannabinoid 2-arachidonoylglycerol mediating inhibition of adenylate cyclase. May function in inflammatory response, nociceptive transmission and bone homeostasis.

**References**

Munro S.,et al.Nature 365:61-65(1993).  
Liu Q.-R.,et al.Genes Brain Behav. 8:519-530(2009).  
Bruess M.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.  
Warren C.N.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.  
Saravanan T.,et al.Submitted (NOV-2005) to the EMBL/GenBank/DDBJ databases.

**Images**

Anti-CNR2 Antibody (N-term) at 1:2000 dilution + human brain lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.