10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

(Mouse) Rnf2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21227c

Product Information

Application	WB, E
Primary Accession	Q9CQJ4
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52402
Calculated MW	37623

Additional Information

Gene ID	19821
Other Names	E3 ubiquitin-protein ligase RING2, 632-, RING finger protein 1B, RING1b, RING finger protein 2, Rnf2, DinG, Ring1b
Target/Specificity	This mouse Rnf2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 165-199 amino acids from the Central region of mouse Rnf2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	(Mouse) Rnf2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Rnf2
Synonyms	DinG, Ring1b
Function	E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and

	gene regulation (PubMed:15525528, PubMed:22325148, PubMed:28596365). H2AK119Ub gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals (PubMed:15525528, PubMed:28596365). May be involved in the initiation of both imprinted and random X inactivation (PubMed:15525528). Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:16710298, PubMed:22325148). PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility (PubMed:15525528, PubMed:16710298, PubMed:22325148). E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4 (PubMed:16710298). Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity (PubMed:15525528, PubMed:16710298). Plays a role in the transcriptional repression of genes that are required for pluripotency in embryonic stem cells, thereby contributing to differentiation of the ectodermal and endodermal germ layers (PubMed:22226355). Association with the chromosomal DNA is cell-cycle dependent. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes (PubMed:24034696). Also acts as a negative regulator of autophagy by mediating ubiquitination of AMBRA1, leading to its subsequent degradation (PubMed:24980959).
Cellular Location	Nucleus. Cytoplasm Chromosome Note=Enriched on inactive X chromosome (Xi) in female trophoblast stem (TS) cells as well as differentiating embryonic stem (ES) cells (PubMed:12183370). The enrichment on Xi is transient during TS and ES cell differentiation. The association with Xi is mitotically stable in non-differentiated TS cells (PubMed:12183370). Co-localizes with SAMD7 in nuclear polycomb bodies (PubMed:28900001)
Tissue Location	Expressed in embryonic stem cells.


Background

E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and gene regulation. H2AK119Ub gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. May be involved in the initiation of both imprinted and random X inactivation. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4. Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity. Association to the chromosomal DNA is cell-cycle dependent. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes.


References

Schoorlemmer J.,et al.EMBO J. 16:5930-5942(1997).
Carninci P.,et al.Science 309:1559-1563(2005).
Ebert L.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Garcia E.,et al.EMBO J. 18:3404-3418(1999).
Suzuki M.,et al.Development 129:4171-4183(2002).

Images



Anti-Rnf2 Antibody (Center) at 1:1000 dilution + F9 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-Rnf2 Antibody (Center) at 1:2000 dilution Lane 1: NCCIT whole cell lysates Lane 2: NIH/3T3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.