

CHST1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21231b

Product Information

Application	WB, IHC-P, E
Primary Accession	O43916
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52426
Calculated MW	46715

Additional Information

Gene ID	8534
Target/Specificity	This CHST1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 392-426 amino acids from the C-terminal region of human CHST1.
Dilution	WB~~1:2000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CHST1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CHST1 (HGNC:1969)
Function	Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of internal galactose (Gal) residues of keratan. Cooperates with B4GALT4 and B3GNT7 glycosyltransferases and CHST6 sulfotransferase to construct and elongate disulfated disaccharide unit [->3(6- sulfoGalbeta)1->4(6-sulfoGlcNAcbeta)1->] within keratan sulfate polymer (PubMed: 10642612 , PubMed: 17690104 , PubMed: 9405439). Has a preference for sulfating keratan sulfate, but it also transfers sulfate to the unsulfated polymer (PubMed: 9405439). Involved in

biosynthesis of phosphacan, a major keratan sulfate proteoglycan in the developing brain (By similarity). Involved in biosynthesis of 6-sulfoGalbeta-containing O-linked glycans in high endothelial venules of lymph nodes. May act in a synergistic manner with CHST4 to generate sialyl 6',6'- disulfo Lewis X motif, a recognition determinant for immune cell receptors implicated in leukocyte trafficking (PubMed:[10330415](#)). Catalyzes sulfation of N-acetyllactosamine (LacNAc) oligosaccharides with highest efficiency for sialylated LacNAc structures (PubMed:[10642612](#)).


Cellular Location

Golgi apparatus membrane; Single- pass type II membrane protein


Tissue Location

Widely expressed at low level. Expressed in brain and skeletal muscle. Expressed by high endothelial cells (HEVs) and leukocytes.

Images



AP21231b staining CHST1 in Human skeletal muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-CHST1 Antibody (C-term) at 1:2000 dilution
 Lane 1: human brain lysates Lane 2: Hela whole cell lysates Lane 3: Jurkat whole cell lysates Lane 4: mouse brain lysates Lane 5: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.