

(Mouse) Ehmt2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21305c

Product Information

Application	WB, E
Primary Accession	Q9Z148
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52371
Calculated MW	138039

Additional Information

Gene ID	110147
Other Names	Histone-lysine N-methyltransferase EHMT2, 211-, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Protein G9a, Ehmt2, Bat8, G9a, Ng36
Target/Specificity	This Mouse Ehmt2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 589-623 amino acids from the Central region of Mouse Ehmt2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	(Mouse) Ehmt2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Ehmt2
Synonyms	Bat8, G9a, Ng36
Function	Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin.

H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q96KQ7}. Chromosome {ECO:0000250|UniProtKB:Q96KQ7}. Note=Almost excluded form nucleoli. Associates with euchromatic regions (By similarity). Does not associate with heterochromatin (By similarity) {ECO:0000250|UniProtKB:Q96KQ7}

Tissue Location

Ubiquitous..


Background

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys- 373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

References

Tachibana M.,et al.Genes Dev. 16:1779-1791(2002).
 Xie T.,et al.Genome Res. 13:2621-2636(2003).
 Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).
 Brown S.E.,et al.Mamm. Genome 12:916-924(2001).
 Tachibana M.,et al.J. Biol. Chem. 276:25309-25317(2001).

Images



Anti-Ehmt2 Antibody (Center)at 1:1000 dilution + mouse brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 138 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.