

# GATA2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21314c

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P23769</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52409
Calculated MW	50500

## Additional Information

---

Gene ID	2624
Other Names	Endothelial transcription factor GATA-2, GATA-binding protein 2, GATA2
Target/Specificity	This GATA2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 212-246 amino acids from the Central region of human GATA2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GATA2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	GATA2
Function	Transcriptional activator which regulates endothelin-1 gene expression in endothelial cells. Binds to the consensus sequence 5'- AGATAG-3'.
Cellular Location	Nucleus.
Tissue Location	Endothelial cells.

## Background

---

Transcriptional activator which regulates endothelin-1 gene expression in endothelial cells. Binds to the consensus sequence 5'-AGATAG-3'.


## References

---

Lee M.-E.,et al.J. Biol. Chem. 266:16188-16192(1991).  
Dorfman D.M.,et al.J. Biol. Chem. 267:1279-1285(1992).  
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Muzny D.M.,et al.Nature 440:1194-1198(2006).

## Images

---



All lanes: Anti-GATA2 Antibody (Center) at 1:1000 dilution + 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 50 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.