

ADAMTS6 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21355a

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, E |
| Primary Accession | Q9UKP5 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB52744 |
| Calculated MW | 125273 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 11174 |
| Other Names | A disintegrin and metalloproteinase with thrombospondin motifs 6, ADAM-TS 6, ADAM-TS6, ADAMTS-6, 3424-, ADAMTS6 |
| Target/Specificity | This ADAMTS6 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 118-151 amino acids from the region of human ADAMTS6. |
| Dilution | WB~1:2000 E~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | ADAMTS6 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|--------------------------|--|
| Name | ADAMTS6 |
| Cellular Location | Secreted, extracellular space, extracellular matrix |
| Tissue Location | Expressed at low levels in placenta and barely detectable in a number of other tissues |

References

Hurskainen T.L.,et al.J. Biol. Chem. 274:25555-25563(1999).


Bevitt D.J.,et al.Gene 359:99-110(2005).

Schmutz J.,et al.Nature 431:268-274(2004).

Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Images



All lanes : Anti-ADAMTS6 Antibody (N-Term) at 1:2000 dilution Lane 1: HeLa whole cell lysates Lane 2: HT-1080 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 125 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.