

# PYCRL Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21368b

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q53H96</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB52797
<b>Calculated MW</b>	28663

## Additional Information

---

<b>Gene ID</b>	65263
<b>Other Names</b>	Pyrroline-5-carboxylate reductase 3, P5C reductase 3, P5CR 3, Pyrroline-5-carboxylate reductase-like protein, PYCRL
<b>Target/Specificity</b>	This PYCRL antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 221-255 amino acids from the C-terminal region of human PYCRL.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PYCRL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PYCR3 ( <a href="#">HGNC:25846</a> )
<b>Function</b>	Oxidoreductase that catalyzes the last step in proline biosynthesis, which corresponds to the reduction of pyrroline-5- carboxylate (P5C) to L-proline using NAD(P)H (PubMed: <a href="#">23024808</a> , PubMed: <a href="#">36414121</a> ). Proline is synthesized from either glutamate or ornithine; both are converted to P5C, and then to proline via pyrroline-5-carboxylate reductases (PYCRs) (PubMed: <a href="#">23024808</a> ).

PYCR3 is exclusively linked to the biosynthesis of proline from ornithine (PubMed:[23024808](#)).

## Cellular Location

Cytoplasm.

## References

---

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Nusbaum C.,et al.Nature 439:331-335(2006).


Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

## Images

---



All lanes : Anti-PYCRL Antibody (C-term) at 1:2000 dilution  
Lane 1: A549 whole cell lysates Lane 2: HeLa whole cell lysates Lane 3: KG-1 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 29 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.