

# KCNMB2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21369a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q9Y691</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB52678

## Additional Information

---

<b>Other Names</b>	Calcium-activated potassium channel subunit beta-2, BK channel subunit beta-2, BKbeta2, Hbeta2, Calcium-activated potassium channel, subfamily M subunit beta-2, Charybdotoxin receptor subunit beta-2, Hbeta3, K(VCA)beta-2, Maxi K channel subunit beta-2, Slo-beta-2, KCNMB2
<b>Target/Specificity</b>	This KCNMB2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-66 amino acids from the N-terminal region of human KCNMB2.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	KCNMB2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

### Background

---

Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel. Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to KCNMA1 channel diversity. Acts as a negative regulator that confers rapid and complete inactivation of KCNMA1 channel complex. May participate in KCNMA1 inactivation in chromaffin cells of the adrenal gland or in hippocampal CA1 neurons.


## References

---

Wallner M., et al. Proc. Natl. Acad. Sci. U.S.A. 96:4137-4142(1999).  
Brenner R., et al. J. Biol. Chem. 275:6453-6461(2000).  
Xia X.-M., et al. J. Neurosci. 19:5255-5264(1999).  
Meera P., et al. Proc. Natl. Acad. Sci. U.S.A. 97:5562-5567(2000).  
Xia X.-M., et al. J. Gen. Physiol. 121:125-148(2003).

## Images

---



All lanes : Anti-KCNMB2 Antibody (N-term) at 1:2000 dilution Lane 1: human brain lysates Lane 2: human heart lysates Lane 3: human kidney lysates Lane 4: HepG2 whole cell lysates Lane 5: mouse brain lysates Lane 6: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.