

HPN Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21383c

Product Information

Application	WB, E
Primary Accession	P05981
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53000
Calculated MW	45011

Additional Information

Gene ID	3249
Other Names	Serine protease hepsin, Transmembrane protease serine 1, Serine protease hepsin non-catalytic chain, Serine protease hepsin catalytic chain, HPN, TMPRSS1
Target/Specificity	This HPN antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 153-188 amino acids from the Central region of human HPN.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HPN Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HPN
Synonyms	TMPRSS1
Function	Serine protease that cleaves extracellular substrates, and contributes to the proteolytic processing of growth factors, such as HGF and MST1/HGFL

(PubMed:[15839837](#), PubMed:[21875933](#)). Plays a role in cell growth and maintenance of cell morphology (PubMed:[21875933](#), PubMed:[8346233](#)). Plays a role in the proteolytic processing of ACE2 (PubMed:[24227843](#)). Mediates the proteolytic cleavage of urinary UMOD that is required for UMOD polymerization (PubMed:[26673890](#)).

Cellular Location

Cell membrane; Single-pass type II membrane protein Apical cell membrane; Single-pass type II membrane protein

Tissue Location

Detected in liver and kidney.


Background

Plays an essential role in cell growth and maintenance of cell morphology. May mediate the activating cleavage of HGF and MST1/HGFL. Plays a role in the proteolytic processing of ACE2.


References

Leytus S.P.,et al.Biochemistry 27:1067-1074(1988).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Tsuji A.,et al.J. Biol. Chem. 266:16948-16953(1991).
Torres-Rosado A.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:7181-7185(1993).
Chen R.,et al.J. Proteome Res. 8:651-661(2009).

Images



All lanes : Anti-HPN Antibody (Center) at 1:2000 dilution
Lane 1: HeLa whole cell lysates Lane 2: SW620 whole cell lysates Lane 3: MCF-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-HPN Antibody (Center)at 1:2000 dilution + mouse liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.