

RFX4 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21391b

Product Information

Application	WB, E
Primary Accession	Q33E94
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52917
Calculated MW	83368

Additional Information

Gene ID	5992
Other Names	Transcription factor RFX4, Regulatory factor X 4, Testis development protein NYD-SP10, RFX4
Target/Specificity	This RFX4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 684-720 amino acids from the C-terminal region of human RFX4.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RFX4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RFX4
Function	Transcription factor that plays a role in early brain development. May activate transcription by interacting directly with the X-box. May activate transcription from CX3CL1 promoter through the X-box during brain development. May be required for neural tube ciliogenesis during embryogenesis (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00858}.

Tissue Location

Isoform 1: Expressed in brain and gliomas (at protein level). Isoform 2: Testis-specific (at protein level). Isoform 3: Testis-specific (at protein level). Isoform 3: Expressed at a higher level in adult testes and ejaculated spermatozoa than in fetal testes Isoform 4: Testis-specific.


Background

May activate transcription by interacting directly with the X-box.


References

Blackshear P.J.,et al.Development 130:4539-4552(2003).
Matsushita H.,et al.Cancer Sci. 96:801-809(2005).
Huang X.,et al.J. Nanosci. Nanotechnol. 5:1236-1239(2005).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Scherer S.E.,et al.Nature 440:346-351(2006).

Images



Anti-RFX4 Antibody (C-term)at 1:2000 dilution + HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-RFX4 Antibody (C-term)at 1:2000 dilution + human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.