10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

p16-INK4A(S152) Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21427a

Product Information

Application	WB, E
Primary Accession	P42771
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53347
Calculated MW	16533

Additional Information

Gene ID	1029
Other Names	Cyclin-dependent kinase inhibitor 2A, isoforms 1/2/3, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A, CDKN2, MTS1
Target/Specificity	This p16-INK4A(S152) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 145-176 amino acids from human p16-INK4A.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	p16-INK4A(S152) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CDKN2A (HGNC:1787)
Synonyms	CDKN2, MTS1
Function	Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact

	with cyclins D and to phosphorylate the retinoblastoma protein.
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific


Background

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

References

Serrano M.,et al.Nature 366:704-707(1993).
Robertson K.D.,et al.Oncogene 18:3810-3820(1999).
Kitagawa Y.,et al.J. Biol. Chem. 277:46289-46297(2002).
Lin Y.C.,et al.Oncogene 26:7017-7027(2007).
Humphray S.J.,et al.Nature 429:369-374(2004).

Images



Anti p16-INK4A(S152) Antibody at 1:2000 dilution + Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 17 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.