

WNT2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21468c

Product Information

Application	WB, E
Primary Accession	P09544
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB49732
Calculated MW	40418

Additional Information

Gene ID	7472
Other Names	Protein Wnt-2, Int-1-like protein 1, Int-1-related protein, IRP, WNT2, INT1L1, IRP
Target/Specificity	This WNT2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 254-287 amino acids from the Central region of human WNT2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WNT2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WNT2
Synonyms	INT1L1, IRP
Function	Ligand for members of the frizzled family of seven transmembrane receptors. Functions in the canonical Wnt signaling pathway that results in activation of transcription factors of the TCF/LEF family (PubMed: 20018874).

Functions as a upstream regulator of FGF10 expression. Plays an important role in embryonic lung development. May contribute to embryonic brain development by regulating the proliferation of dopaminergic precursors and neurons (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted

Tissue Location

Expressed in brain in the thalamus, in fetal and adult lung and in placenta.


Background

Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule which affects the development of discrete regions of tissues. Is likely to signal over only few cell diameters.

References

Wainwright B.J.,et al.EMBO J. 7:1743-1748(1988).
Farrall M.,et al.Submitted (APR-1988) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Hillier L.W.,et al.Nature 424:157-164(2003).

Images



All lanes : Anti-WNT2 Antibody (Center) at 1:2000 dilution
Lane 1: Jurkat whole cell lysates Lane 2: human placenta lysates Lane 3: human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.