

## PICALM Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21487c

### Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q13492</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB53756
<b>Calculated MW</b>	70755

### Additional Information

---

<b>Gene ID</b>	8301
<b>Other Names</b>	Phosphatidylinositol-binding clathrin assembly protein, Clathrin assembly lymphoid myeloid leukemia protein, PICALM, CALM
<b>Target/Specificity</b>	This PICALM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 385-415 amino acids from the Central region of human PICALM.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PICALM Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

---

<b>Name</b>	PICALM
<b>Synonyms</b>	CALM
<b>Function</b>	Cytoplasmic adapter protein that plays a critical role in clathrin-mediated endocytosis which is important in processes such as internalization of cell receptors, synaptic transmission or removal of apoptotic cells. Recruits AP-2

and attaches clathrin triskelions to the cytoplasmic side of plasma membrane leading to clathrin-coated vesicles (CCVs) assembly (PubMed:[10436022](#), PubMed:[16262731](#), PubMed:[27574975](#)). Furthermore, regulates clathrin-coated vesicle size and maturation by directly sensing and driving membrane curvature (PubMed:[25898166](#)). In addition to binding to clathrin, mediates the endocytosis of small R- SNARES (Soluble NSF Attachment Protein REceptors) between plasma membranes and endosomes including VAMP2, VAMP3, VAMP4, VAMP7 or VAMP8 (PubMed:[21808019](#), PubMed:[22118466](#), PubMed:[23741335](#)). In turn, PICALM- dependent SNARE endocytosis is required for the formation and maturation of autophagic precursors (PubMed:[25241929](#)). Modulates thereby autophagy and the turnover of autophagy substrates such as MAPT/TAU or amyloid precursor protein cleaved C-terminal fragment (APP- CTF) (PubMed:[24067654](#), PubMed:[25241929](#)).

#### Cellular Location

Cell membrane. Membrane, clathrin-coated pit. Golgi apparatus. Cytoplasmic vesicle, clathrin- coated vesicle. Nucleus. Note=Colocalized with clathrin in the Golgi area (PubMed:10436022). Interaction with PIMREG may target PICALM to the nucleus in some cells (PubMed:16491119)

#### Tissue Location

Expressed in all tissues examined.


## Background

Assembly protein recruiting clathrin and adapter protein complex 2 (AP2) to cell membranes at sites of coated-pit formation and clathrin-vesicle assembly. May be required to determine the amount of membrane to be recycled, possibly by regulating the size of the clathrin cage. Involved in AP2-dependent clathrin-mediated endocytosis at the neuromuscular junction.

## References

Dreyling M.H.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:4804-4809(1996).  
 Ota T.,et al.Nat. Genet. 36:40-45(2004).  
 Nakajima D.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.  
 Taylor T.D.,et al.Nature 440:497-500(2006).  
 Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

## Images



All lanes : Anti-PICALM Antibody (Center) at 1:2000 dilution Lane 1: A431 whole cell lysates Lane 2: A549 whole cell lysates Lane 3: HeLa whole cell lysates Lane 4: NIH/3T3 whole cell lysates Lane 5: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.