10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# PGM1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21492b

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, E |
| **Primary Accession** | P36871 |
| **Reactivity** | Human, Rat, Mouse |
| **Predicted** | Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB53795 |
| **Calculated MW** | 61449 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 5236 |
| **Other Names** | Phosphoglucomutase-1, PGM 1, Glucose phosphomutase 1, PGM1 |
| **Target/Specificity** | This PGM1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 469-501 amino acids from human PGM1. |
| **Dilution** | WB~~1:2000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | PGM1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | PGM1 |
| **Function** | Catalyzes the reversible isomerization of alpha-D-glucose 1- phosphate to alpha-D-glucose 6-phosphate (PubMed:15378030, PubMed:25288802). The mechanism proceeds via the intermediate compound alpha-D-glucose 1,6-bisphosphate (Probable) (PubMed:25288802). This enzyme participates in both the breakdown and synthesis of glucose (PubMed:17924679, |

PubMed:25288802).

**Cellular Location** [Isoform 1]: Cytoplasm.


## Background

This enzyme participates in both the breakdown and synthesis of glucose.


## References

Whitehouse D.B.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:411-415(1992).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Putt W.,et al.Biochem. J. 296:417-422(1993).

## Images



AP21492b staining PGM1 in human heart tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.




All lanes : Anti-PGM1 Antibody (C-Term) at 1:2000 dilution Lane 1: mouse brain lysates Lane 2: mouse heart lysates Lane 3: human liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.