10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HOOK2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21495a

Product Information

Application	WB, E
Primary Accession	Q96ED9
Reactivity	Human, Mouse
Predicted	Rat, Bovine, Canine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB49361
Calculated MW	83207

Additional Information

Gene ID	29911
Other Names	Protein Hook homolog 2, h-hook2, hHK2, HOOK2
Target/Specificity	This HOOK2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 162-197 amino acids from the N-terminal region of human HOOK2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HOOK2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HOOK2
Function	Component of the FTS/Hook/FHIP complex (FHF complex). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex (the HOPS complex). Contributes to the establishment and maintenance of centrosome function. May function in the positioning or formation of aggresomes, which are pericentriolar

accumulations of misfolded proteins, proteasomes and chaperones. FHF complex promotes the distribution of AP-4 complex to the perinuclear area of the cell (PubMed:32073997).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q7TMK6}. Golgi apparatus, trans-Golgi network. Note=Colocalizes with aggresomes, which are aggregates of misfolded proteins, at the centrosome (PubMed:17540036). Also localizes to punctate cytoplasmic foci which do not appear to overlap with early or late endosomes, the endoplasmic reticulum, multivesicular bodies (MVBs), lysosome, or mitochondria (PubMed:17540036, PubMed:32073997). Often found in close association with microtubules (PubMed:17540036). Localizes to the manchette in elongating spermatids (By similarity). {ECO:0000250|UniProtKB:Q7TMK6, ECO:0000269|PubMed:17540036, ECO:0000269|PubMed:32073997}


Background

Component of the FTS/Hook/FHIP complex (FHF complex). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex (the HOPS complex). Contributes to the establishment and maintenance of centrosome function. May function in the positioning or formation of aggresomes, which are pericentriolar accumulations of misfolded proteins, proteasomes and chaperones.

References

Kraemer H.,et al.Genetics 151:675-684(1999).
Grimwood J.,et al.Nature 428:529-535(2004).
Walenta J.H.,et al.J. Cell Biol. 152:923-934(2001).
Szebenyi G.,et al.BMC Cell Biol. 8:19-19(2007).
Szebenyi G.,et al.Traffic 8:32-46(2007).

Images



Anti-HOOK2 Antibody (N-term)at 1:2000 dilution + MCF-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.