

AKR7L Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21497a

Product Information

Application	WB, IF, E
Primary Accession	Q8NHP1
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53759
Calculated MW	36970

Additional Information

Gene ID	246181
Other Names	Aflatoxin B1 aldehyde reductase member 4, 1---, AFB1 aldehyde reductase 3, AFB1-AR 3, Aldoketoreductase 7-like, AKR7L, AFAR3, AKR7A4
Target/Specificity	This AKR7L antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-64 amino acids from human AKR7L.
Dilution	WB~~1:2000 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	AKR7L Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AKR7L
Synonyms	AFAR3 {ECO:0000303 PubMed:12879023}, AKR
Function	Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

Tissue Location

Mainly expressed in uterus.

Background


Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen (By similarity).

References


Gregory S.G.,et al.Nature 441:315-321(2006).

Praml C.,et al.Oncogene 22:4765-4773(2003).

Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HepG2 (human liver hepatocellular carcinoma cell line) cells labeling AKR7L with AP21497a at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on HepG2 cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).



Anti-AKR7L Antibody (N-Term)at 1:2000 dilution + HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.