

GSTT1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21534b

Product Information

Application	WB, E
Primary Accession	P30711
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB47964
Calculated MW	27335

Additional Information

Gene ID	2952
Other Names	Glutathione S-transferase theta-1, GST class-theta-1, Glutathione transferase T1-1, GSTT1
Target/Specificity	This GSTT1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 216-250 amino acids from the C-terminal region of human GSTT1.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GSTT1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GSTT1
Function	Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Acts on 1,2-epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with cumene hydroperoxide.

Cellular Location	Cytoplasm.
Tissue Location	Found in erythrocyte. Expressed at low levels in liver. In lung, expressed at low levels in club cells and ciliated cells at the alveolar/bronchiolar junction. Absent from epithelial cells of larger bronchioles.


Background

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Acts on 1,2- epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4- nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with cumene hydroperoxide.

References

Pemble S.,et al.Biochem. J. 300:271-276(1994).
Jemth P.,et al.Arch. Biochem. Biophys. 348:247-254(1997).
Sprenger R.,et al.Pharmacogenetics 10:557-565(2000).
Iida A.,et al.Submitted (MAR-2001) to the EMBL/GenBank/DDBJ databases.
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).

Images



All lanes : Anti-GSTT1 Antibody (C-term) at 1:2000 dilution
Lane 1: Hela whole cell lysates Lane 2: HepG2 whole cell lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 27 kDa
Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.