

RLF Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21542b

Product Information

Application	WB, E
Primary Accession	Q13129
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53780
Calculated MW	217953

Additional Information

Gene ID	6018
Other Names	Zinc finger protein Rlf, Rearranged L-myc fusion gene protein, Zn-15-related protein, RLF
Target/Specificity	This RLF antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1751-1785 amino acids from human RLF.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RLF Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RLF
Function	May be involved in transcriptional regulation.
Cellular Location	Nucleus.

Tissue Location

Widely expressed in fetal and adult tissues.


Background

May be involved in transcriptional regulation.

References

Makela T.P.,et al.Oncogene 11:2699-2704(1995).
Gregory S.G.,et al.Nature 441:315-321(2006).
Olsen J.V.,et al.Cell 127:635-648(2006).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).
Rigbolt K.T.,et al.Sci. Signal. 4:RS3-RS3(2011).

Images



Anti-RLF Antibody (C-Term)at 1:2000 dilution + PC-3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 218 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.