

# MTCP1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21543b

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P56278</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB53779
<b>Calculated MW</b>	12600

## Additional Information

---

<b>Gene ID</b>	4515
<b>Other Names</b>	Protein p13 MTCP-1, p13MTCP1, Mature T-cell proliferation-1 type B1, MTCP-1 type B1, MTCP1, C61B
<b>Target/Specificity</b>	This MTCP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 73-103 amino acids from human MTCP1.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MTCP1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MTCP1
<b>Synonyms</b>	C6.1B
<b>Function</b>	Enhances the phosphorylation and activation of AKT1 and AKT2.
<b>Tissue Location</b>	Not found at a significant level in any tissue.

## Background

---

Enhances the phosphorylation and activation of AKT1 and AKT2.

## References

---

Stern M.-H.,et al.Oncogene 8:2475-2483(1993).

Ross M.T.,et al.Nature 434:325-337(2005).


Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Laine J.,et al.Mol. Cell 6:395-407(2000).

Laine J.,et al.J. Biol. Chem. 277:3743-3751(2002).

## Images

---



All lanes : Anti-MTCP1 Antibody (C-Term) at 1:2000 dilution Lane 1: mouse kidney lysates Lane 2: human liver lysates Lane 3: human kidney lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 13 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.