

OTUB1 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21558b

Product Information

Application	WB, E
Primary Accession	<u>Q96FW1</u>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53068
Calculated MW	31284

Additional Information

Gene ID	55611
Other Names	Ubiquitin thioesterase OTUB1, Deubiquitinating enzyme OTUB1, OTU domain-containing ubiquitin aldehyde-binding protein 1, Otubain-1, hOTU1, Ubiquitin-specific-processing protease OTUB1, OTUB1, OTB1, OTU1
Target/Specificity	This OTUB1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 185-219 amino acids from human OTUB1.
Dilution	WB~~1:1000-1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	OTUB1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	OTUB1
Synonyms	OTB1, OTU1
Function	Hydrolase that can specifically remove 'Lys-48'-linked conjugated ubiquitin from proteins and plays an important regulatory role at the level of protein

	turnover by preventing degradation (PubMed: <u>12401499</u> , PubMed: <u>12704427</u> ,
	PubMed: <u>14661020</u> , PubMed: <u>23827681</u>). Regulator of T-cell anergy, a
	phenomenon that occurs when T-cells are rendered unresponsive to antigen
	rechallenge and no longer respond to their cognate antigen
	(PubMed: <u>14661020</u>). Acts via its interaction with RNF128/GRAIL, a crucial
	inductor of CD4 T-cell anergy (PubMed: <u>14661020</u>). Isoform 1 destabilizes
	RNF128, leading to prevent anergy (PubMed: <u>14661020</u>). In contrast, isoform 2
	stabilizes RNF128 and promotes anergy (PubMed: <u>14661020</u>). Surprisingly, it
	regulates RNF128- mediated ubiquitination, but does not deubiquitinate
	polyubiquitinated RNF128 (PubMed: <u>14661020</u>). Deubiquitinates estrogen
	receptor alpha (ESR1) (PubMed: <u>19383985</u>). Mediates deubiquitination of
	'Lys-48'-linked polyubiquitin chains, but not 'Lys-63'-linked polyubiquitin
	chains (PubMed: <u>18954305</u> , PubMed: <u>19211026</u> , PubMed: <u>23827681</u>). Not able
	to cleave di-ubiquitin (PubMed: <u>18954305</u> , PubMed: <u>23827681</u>). Also capable of
	removing NEDD8 from NEDD8 conjugates, but with a much lower preference
	compared to 'Lys-48'-linked ubiquitin (PubMed: <u>18954305</u> ,
	PubMed: <u>23827681</u>).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:B2RYG6}.
Tissue Location	Isoform 1 is ubiquitous. Isoform 2 is expressed only in lymphoid tissues such as tonsils, lymph nodes and spleen, as well as peripheral blood mononuclear cells

Background

Hydrolase that can specifically remove 'Lys-48'-linked conjugated ubiquitin from proteins and plays an important regulatory role at the level of protein turnover by preventing degradation. Regulator of T-cell anergy, a phenomenon that occurs when T-cells are rendered unresponsive to antigen rechallenge and no longer respond to their cognate antigen. Acts via its interaction with RNF128/GRAIL, a crucial inductor of CD4 T-cell anergy. Isoform 1 destabilizes RNF128, leading to prevent anergy. In contrast, isoform 2 stabilizes RNF128 and promotes anergy. Surprisingly, it regulates RNF128-mediated ubiquitination, but does not deubiquitinate polyubiquitinated RNF128. Deubiquitinates estrogen receptor alpha (ESR1). Mediates deubiquitination of 'Lys- 48'-linked polyubiquitin chains, but not 'Lys-63'-linked polyubiquitin chains. Not able to cleave di-ubiquitin. Also capable of removing NEDD8 from NEDD8 conjugates, but with a much lower preference compared to 'Lys-48'-linked ubiquitin.

References

Balakirev M.Y.,et al.EMBO Rep. 4:517-522(2003). Soares L.,et al.Nat. Immunol. 5:45-54(2004). Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000). Ota T.,et al.Nat. Genet. 36:40-45(2004). Taylor T.D.,et al.Nature 440:497-500(2006).

Images

All lanes : Anti-OTUB1 Antibody (C-Term) at 1:1000-1:2000 dilution Lane 1: A549 whole cell lysates Lane 2: Hela whole cell lysates Lane 3: HepG2 whole cell lysates Lane 4: MCF-7 whole cell lysates Lane 5: NIH/3T3 whole cell lysates Lane 6: U-2OS whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5%



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.