

# NDUFB7 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21560a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P17568</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB49277
<b>Calculated MW</b>	16402

## Additional Information

---

<b>Gene ID</b>	4713
<b>Other Names</b>	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7, Cell adhesion protein SQM1, Complex I-B18, CI-B18, NADH-ubiquinone oxidoreductase B18 subunit, NDUFB7
<b>Target/Specificity</b>	This NDUFB7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 27-60 amino acids from the N-terminal region of human NDUFB7.
<b>Dilution</b>	WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	NDUFB7 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NDUFB7
<b>Function</b>	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be

ubiquinone.

#### Cellular Location

Mitochondrion inner membrane; Peripheral membrane protein.  
Mitochondrion intermembrane space

## Background

---

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.


## References

---

Wong Y.-C.,et al.Biochem. Biophys. Res. Commun. 166:984-992(1990).  
Triepels R.,et al.Hum. Genet. 106:385-391(2000).  
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).  
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---



Anti-NDUFB7 Antibody (N-term)at 1:2000 dilution +  
mouse ovary lysates Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase  
conjugated at 1/10000 dilution. Predicted band size : 16  
kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.