10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# TRRAP Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21568b

## Product Information

| | |
|---|---|
| **Application** | WB, IF, E |
| **Primary Accession** | Q9Y4A5 |
| **Reactivity** | Human, Mouse |
| **Predicted** | Rat, Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB51386 |
| **Calculated MW** | 437600 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 8295 |
| **Other Names** | Transformation/transcription domain-associated protein, 350/400 kDa PCAF-associated factor, PAF350/400, STAF40, Tra1 homolog, TRRAP, PAF400 |
| **Target/Specificity** | This TRRAP antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 3015-3049 amino acids from the C-terminal region of human TRRAP. |
| **Dilution** | WB~~1:2000 IF~~1:25 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | TRRAP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | TRRAP |
| **Synonyms** | PAF400 |
| **Function** | Adapter protein, which is found in various multiprotein chromatin complexes with histone acetyltransferase activity (HAT), which gives a specific |

tag for epigenetic transcription activation. Component of the NuA4 histone acetyltransferase complex which is responsible for acetylation of nucleosomal histones H4 and H2A. Plays a central role in MYC transcription activation, and also participates in cell transformation by MYC. Required for p53/TP53-, E2F1- and E2F4- mediated transcription activation. Also involved in transcription activation mediated by the adenovirus E1A, a viral oncoprotein that deregulates transcription of key genes. Probably acts by linking transcription factors such as E1A, MYC or E2F1 to HAT complexes such as STAGA thereby allowing transcription activation. Probably not required in the steps following histone acetylation in processes of transcription activation. May be required for the mitotic checkpoint and normal cell cycle progression. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome. May play a role in the formation and maintenance of the auditory system (By similarity).

**Cellular Location** Nucleus


## Background

Adapter protein, which is found in various multiprotein chromatin complexes with histone acetyltransferase activity (HAT), which gives a specific tag for epigenetic transcription activation. Component of the NuA4 histone acetyltransferase complex which is responsible for acetylation of nucleosomal histones H4 and H2A. Plays a central role in MYC transcription activation, and also participates in cell transformation by MYC. Required for p53/TP53-, E2F1- and E2F4-mediated transcription activation. Also involved in transcription activation mediated by the adenovirus E1A, a viral oncoprotein that deregulates transcription of key genes. Probably acts by linking transcription factors such as E1A, MYC or E2F1 to HAT complexes such as STAGA thereby allowing transcription activation. Probably not required in the steps following histone acetylation in processes of transcription activation. May be required for the mitotic checkpoint and normal cell cycle progression. Component of a SWR1- like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.

## References


McMahon S.B.,et al.Cell 94:363-374(1998).
Vassilev A.,et al.Mol. Cell 2:869-875(1998).
Hillier L.W.,et al.Nature 424:157-164(2003).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Scherer S.W.,et al.Science 300:767-772(2003).

## Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling TRRAP with AP21568b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing nucleus staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).

All lanes : Anti-TRRAP Antibody (C-term) at 1:2000 dilution




Lane 1: K562 whole cell lysates Lane 2: Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 438 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.