

Zebrafish ak2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21602c

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q1L8L9
Reactivity	Human, Zebrafish, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53214
Calculated MW	26616

Additional Information

Gene ID	321793
Other Names	Adenylate kinase 2, mitochondrial {ECO:0000255 HAMAP-Rule:MF_03168}, AK 2 {ECO:0000255 HAMAP-Rule:MF_03168}, 2743 {ECO:0000255 HAMAP-Rule:MF_03168}, ATP-AMP transphosphorylase 2 {ECO:0000255 HAMAP-Rule:MF_03168}, ATP:AMP phosphotransferase {ECO:0000255 HAMAP-Rule:MF_03168}, Adenylate monophosphate kinase {ECO:0000255 HAMAP-Rule:MF_03168}, ak2
Target/Specificity	This Zebrafish ak2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 147-183 amino acids from the Central region of Zebrafish ak2.
Dilution	WB~~1:2000 IHC-P~~1:100~500 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Zebrafish ak2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ak2
------	-----

Function	Catalyzes the reversible transfer of the terminal phosphate group between ATP and AMP. Plays an important role in cellular energy homeostasis and in adenine nucleotide metabolism. Adenylate kinase activity is critical for regulation of the phosphate utilization and the AMP de novo biosynthesis pathways. Plays a key role in hematopoiesis.
Cellular Location	Mitochondrion intermembrane space {ECO:0000255 HAMAP-Rule:MF_03168}


Background

Catalyzes the reversible transfer of the terminal phosphate group between ATP and AMP. Plays an important role in cellular energy homeostasis and in adenine nucleotide metabolism. Adenylate kinase activity is critical for regulation of the phosphate utilization and the AMP de novo biosynthesis pathways. Plays a key role in hematopoiesis.


References

Howe K.,et al.Nature 496:498-503(2013).
Pannicke U.,et al.Nat. Genet. 41:101-105(2009).

Images




Overlay histogram showing ZF4 cells stained with AP21602c (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP21602c, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.




AP21602c staining Zebrafish ak2 in human heart tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

AP21602c staining Zebrafish ak2 in zebra fish body tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6).



Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-Zebrafish ak2 Antibody (Center) at 1:2000 dilution Lane 1: Zebrafish lysates Lane 2: ZF4 whole cell lysates Lane 3: human heart lysates Lane 4: mouse liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.