

NGEF Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21608a

Product Information

Application	WB, E
Primary Accession	Q8N5V2
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53236
Calculated MW	82496

Additional Information

Gene ID	25791
Other Names	Ephexin-1, Eph-interacting exchange protein, Neuronal guanine nucleotide exchange factor, NGEF
Target/Specificity	This NGEF antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 187-220 amino acids from the N-terminal region of human NGEF.
Dilution	WB~~1:1000-1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NGEF Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NGEF
Function	Acts as a guanine nucleotide exchange factor (GEF) which differentially activates the GTPases RHOA, RAC1 and CDC42. Plays a role in axon guidance regulating ephrin-induced growth cone collapse and dendritic spine morphogenesis. Upon activation by ephrin through EPHA4, the GEF activity switches toward RHOA resulting in its activation. Activated RHOA promotes

cone retraction at the expense of RAC1- and CDC42-stimulated growth cone extension (By similarity).

Cellular Location

Cytoplasm. Membrane. Cell projection, growth cone. Note=Associated with membranes. Localizes to axonal growth cones (By similarity)

Tissue Location

Highly expressed in brain specifically in caudate nucleus and to a lower extent in amygdala and hippocampus. Also detected in lung.


Background

Acts as a guanine nucleotide exchange factor (GEF) which differentially activates the GTPases RHOA, RAC1 and CDC42. Plays a role in axon guidance regulating ephrin-induced growth cone collapse and dendritic spine morphogenesis. Upon activation by ephrin through EPHA4, the GEF activity switches toward RHOA resulting in its activation. Activated RHOA promotes cone retraction at the expense of RAC1- and CDC42-stimulated growth cone extension (By similarity).

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Rodrigues N.R.,et al.Genomics 65:53-61(2000).

Images



All lanes : Anti-NGEF Antibody (N-term) at 1:1000-1:2000 dilution
Lane 1: A431 whole cell lysates
Lane 2: rat brain lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 82 kDa
Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.