

# GCKR Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21610a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q14397</a>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53063
Calculated MW	68685

## Additional Information

---

Gene ID	2646
Other Names	Glucokinase regulatory protein, GKRP, Glucokinase regulator, GCKR
Target/Specificity	This GCKR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 87-120 amino acids from human GCKR.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GCKR Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	GCKR {ECO:0000303 PubMed:8589523, ECO:0000312 HGNC:HGNC:4196}
Function	Regulates glucokinase (GCK) by forming an inactive complex with this enzyme (PubMed: <a href="#">23621087</a> , PubMed: <a href="#">23733961</a> ). Acts by promoting GCK recruitment to the nucleus, possibly to provide a reserve of GCK that can be quickly released in the cytoplasm after a meal (PubMed: <a href="#">10456334</a> ). The affinity of GCKR for GCK is modulated by fructose metabolites: GCKR with bound fructose 6-phosphate has increased affinity for GCK, while GCKR with bound fructose 1-phosphate has strongly decreased affinity for GCK and does

not inhibit GCK activity (PubMed:[23621087](#), PubMed:[23733961](#)).

#### Cellular Location

Cytoplasm. Nucleus. Mitochondrion {ECO:0000250|UniProtKB:Q07071}.  
Note=Under low glucose concentrations, GCKR associates with GCK and the inactive complex is recruited to the hepatocyte nucleus.

#### Tissue Location

Found in liver and pancreas. Not detected in muscle, brain, heart, thymus, intestine, uterus, adipose tissue, kidney, adrenal, lung or spleen.


## Background

Inhibits glucokinase (GCK) by forming an inactive complex with this enzyme. The affinity of GCKR for GCK is modulated by fructose metabolites: GCKR with bound fructose 6- phosphate has increased affinity for GCK, while GCKR with bound fructose 1-phosphate has strongly decreased affinity for GCK and does not inhibit GCK activity.

## References

Warner J.P.,et al.Mamm. Genome 6:532-536(1995).  
Hayward B.E.,et al.Genomics 49:137-142(1998).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
de la Iglesia N.,et al.FEBS Lett. 456:332-338(1999).

## Images



All lanes : Anti-GCKR Antibody (N-Term) at 1:2000 dilution  
Lane 1: mouse liver lysates Lane 2: rat liver lysates  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.