

GCKR Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21610a

Product Information

Application	WB, E
Primary Accession	Q14397
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53063
Calculated MW	68685

Additional Information

Gene ID	2646
Other Names	Glucokinase regulatory protein, GKRP, Glucokinase regulator, GCKR
Target/Specificity	This GCKR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 87-120 amino acids from human GCKR.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GCKR Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GCKR {ECO:0000303 PubMed:8589523, ECO:0000312 HGNC:HGNC:4196}
Function	Regulates glucokinase (GCK) by forming an inactive complex with this enzyme (PubMed: 23621087 , PubMed: 23733961). Acts by promoting GCK recruitment to the nucleus, possibly to provide a reserve of GCK that can be quickly released in the cytoplasm after a meal (PubMed: 10456334). The affinity of GCKR for GCK is modulated by fructose metabolites: GCKR with bound fructose 6-phosphate has increased affinity for GCK, while GCKR with bound fructose 1-phosphate has strongly decreased affinity for GCK and does

not inhibit GCK activity (PubMed:[23621087](#), PubMed:[23733961](#)).

Cellular Location

Cytoplasm. Nucleus. Mitochondrion {ECO:0000250|UniProtKB:Q07071}.
Note=Under low glucose concentrations, GCKR associates with GCK and the inactive complex is recruited to the hepatocyte nucleus.

Tissue Location

Found in liver and pancreas. Not detected in muscle, brain, heart, thymus, intestine, uterus, adipose tissue, kidney, adrenal, lung or spleen.


Background

Inhibits glucokinase (GCK) by forming an inactive complex with this enzyme. The affinity of GCKR for GCK is modulated by fructose metabolites: GCKR with bound fructose 6- phosphate has increased affinity for GCK, while GCKR with bound fructose 1-phosphate has strongly decreased affinity for GCK and does not inhibit GCK activity.

References

Warner J.P.,et al.Mamm. Genome 6:532-536(1995).
Hayward B.E.,et al.Genomics 49:137-142(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
de la Iglesia N.,et al.FEBS Lett. 456:332-338(1999).

Images



All lanes : Anti-GCKR Antibody (N-Term) at 1:2000 dilution
Lane 1: mouse liver lysates Lane 2: rat liver lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.