

# DSTYK Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21613a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q6XUX3</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53052
Calculated MW	105206

## Additional Information

---

Gene ID	25778
Other Names	Dual serine/threonine and tyrosine protein kinase, Dusty protein kinase, Dusty PK, RIP-homologous kinase, Receptor-interacting serine/threonine-protein kinase 5, Sugen kinase 496, Sgk496, DSTYK, KIAA0472, RIP5, RIPK5, SGK496
Target/Specificity	This DSTYK antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 256-290 amino acids from human DSTYK.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DSTYK Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	DSTYK
Synonyms	KIAA0472, RIP5, RIPK5, SGK496
Function	Acts as a positive regulator of ERK phosphorylation downstream of

fibroblast growth factor-receptor activation (PubMed:[23862974](#), PubMed:[28157540](#)). Involved in the regulation of both caspase-dependent apoptosis and caspase-independent cell death (PubMed:[15178406](#)). In the skin, it plays a predominant role in suppressing caspase-dependent apoptosis in response to UV stress in a range of dermal cell types (PubMed:[28157540](#)).

#### Cellular Location

Cytoplasm. Cell membrane {ECO:0000250|UniProtKB:Q6XUX1}. Apical cell membrane. Basolateral cell membrane. Cell junction {ECO:0000250|UniProtKB:Q6XUX1}. Note=Detected at apical cell-cell junctions. Colocalized with FGF receptors to the cell membrane (By similarity). Detected in basolateral and apical membranes of all tubular epithelia. {ECO:0000250|UniProtKB:Q6XUX1, ECO:0000269|PubMed:23862974}

#### Tissue Location

Predominantly expressed in skeletal muscle and testis. Expressed in basolateral and apical membranes of all tubular epithelia. Expressed in thin ascending limb of the loop of Henle and the distal convoluted tubule. Expressed in all layers of transitional ureteric epithelium and in the ureteric smooth-muscle cells. Weakly expressed in heart, brain, placenta, kidney, pancreas, spleen, thymus, prostate, uterus, small intestine, white blood cells, stomach, spinal cord and adrenal gland. Is widely distributed in the CNS. Also detected in several tumor cell lines. Expressed in the skin (PubMed:28157540)


## Background

Acts as a positive regulator of ERK phosphorylation downstream of fibroblast growth factor-receptor activation. May induce both caspase-dependent apoptosis and caspase-independent cell death.

## References

Peng J.,et al.Biochim. Biophys. Acta 1759:562-572(2006).  
Zhao Z.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.  
Gregory S.G.,et al.Nature 441:315-321(2006).  
Seki N.,et al.DNA Res. 4:345-349(1997).  
Zha J.,et al.Biochem. Biophys. Res. Commun. 319:298-303(2004).

## Images



Anti-DSTYK Antibody (N-Term)at 1:2000 dilution + rat brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 105 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.