

# PRSS21 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21624a

#### **Product Information**

**Application** WB, E **Primary Accession Q9Y6M0** Reactivity Human Host Rabbit Clonality polyclonal Isotype Rabbit IgG **Clone Names** RB53141 Calculated MW 34884

#### **Additional Information**

**Gene ID** 10942

Other Names Testisin, 3421-, Eosinophil serine protease 1, ESP-1, Serine protease 21,

PRSS21, ESP1, TEST1

**Target/Specificity** This PRSS21 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 14-26 amino acids from the N-Term

region of human PRSS21.

**Dilution** WB~~1:1000-1:2000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** PRSS21 Antibody (N-Term) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name PRSS21

Synonyms ESP1, TEST1

**Function** Could regulate proteolytic events associated with testicular germ cell

maturation.

**Cellular Location** Cell membrane; Lipid-anchor, GPI- anchor

**Tissue Location** Expressed predominantly in premeiotic testicular germ cells, mostly late

pachytene and diplotene spermatocytes


## **Background**

Could regulate proteolytic events associated with testicular germ cell maturation.

#### References

Inoue M., et al. Biochem. Biophys. Res. Commun. 252:307-312(1998). Inoue M., et al. Biochem. Biophys. Res. Commun. 266:564-568(1999). Hooper J.D., et al. Cancer Res. 59:3199-3205(1999). Hooper J.D., et al. Biochim. Biophys. Acta 1492:63-71(2000). Clark H.F., et al. Genome Res. 13:2265-2270(2003).

### **Images**



All lanes: Anti-PRSS21 Antibody (N-Term) at 1:1000-1:2000 dilution Lane 1: K562 whole cell lysates Lane 2: Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 35 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.