

PSMB9 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21630b

Product Information

Application	WB, E
Primary Accession	P28065
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53298

Additional Information

Other Names	Proteasome subunit beta type-9, Low molecular mass protein 2, Macropain chain 7, Multicatalytic endopeptidase complex chain 7, Proteasome chain 7, Proteasome subunit beta-1i, Really interesting new gene 12 protein, PSMB9, LMP2, PSMB6i, RING12
Target/Specificity	This PSMB9 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 189-219 amino acids from the human region of human PSMB9.
Dilution	WB~1:2000 E~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PSMB9 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Background

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. This subunit is involved in antigen processing to generate class I binding peptides. Replacement of PSMB6 by PSMB9 increases the capacity of the immunoproteasome to cleave model peptides after hydrophobic and basic residues.

References

Glynn R., et al. Eur. J. Immunol. 23:860-866(1993).
Beck S., et al. J. Mol. Biol. 228:433-441(1992).
Kelly A., et al. Nature 353:667-668(1991).
Fruh K., et al. J. Biol. Chem. 267:22131-22140(1992).
Beck S., et al. J. Mol. Biol. 255:1-13(1996).

Images



All lanes : Anti-PSMB9 Antibody (CTerm) at 1:2000 dilution
Lane 1: rat testis lysate Lane 2: U-2OS whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.