

WIPF3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21633a

Product Information

ApplicationWB, EPrimary AccessionA6NGB9

Reactivity Human, Rat, Mouse

Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Clone Names RB43359
Calculated MW 49458

Additional Information

Gene ID 644150

Other Names WAS/WASL-interacting protein family member 3, Corticosteroids and regional

expression protein 16 homolog, WIPF3, CR16

Target/Specificity This WIPF3 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 109-142 amino acids from the

N-terminal region of human WIPF3.

Dilution WB~~1:2000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions WIPF3 Antibody (N-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name WIPF3

Synonyms CR16

Function May be a regulator of cytoskeletal organization. May have a role in

spermatogenesis (By similarity).

Cytoplasm. Note=In hippocampal neurons colocalizes with WASL in the cell body, axons and the growth cone


Background

May be a regulator of cytoskeletal organization. May have a role in spermatogenesis (By similarity).

References

Hillier L.W.,et al.Nature 424:157-164(2003). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Ebert L.,et al.Submitted (JAN-2003) to the EMBL/GenBank/DDBJ databases. Zettl M.,et al.Curr. Biol. 12:1617-1622(2002).

Images



All lanes: Anti-WIPF3 Antibody (N-term) at 1:2000 dilution Lane 1: human liver lysate Lane 2: rat liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 49 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.