

ABCB11 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP21646b

Product Information

Application	WB, E
Primary Accession	O95342
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB53271
Calculated MW	146407

Additional Information

Gene ID	8647
Other Names	Bile salt export pump, ATP-binding cassette sub-family B member 11, ABCB11, BSEP
Target/Specificity	This ABCB11 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1063-1097 amino acids of human ABCB11.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ABCB11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ABCB11 (HGNC:42)
Synonyms	BSEP {ECO:0000303 Ref.2}
Function	Catalyzes the transport of the major hydrophobic bile salts, such as taurine and glycine-conjugated cholic acid across the canalicular membrane of hepatocytes in an ATP-dependent manner, therefore participates in hepatic

bile acid homeostasis and consequently to lipid homeostasis through regulation of biliary lipid secretion in a bile salts dependent manner (PubMed:[15791618](#), PubMed:[16332456](#), PubMed:[18985798](#), PubMed:[19228692](#), PubMed:[20010382](#), PubMed:[20398791](#), PubMed:[22262466](#), PubMed:[24711118](#), PubMed:[29507376](#), PubMed:[32203132](#)). Transports taurine-conjugated bile salts more rapidly than glycine-conjugated bile salts (PubMed:[16332456](#)). Also transports non-bile acid compounds, such as pravastatin and fexofenadine in an ATP-dependent manner and may be involved in their biliary excretion (PubMed:[15901796](#), PubMed:[18245269](#)).

Cellular Location

Apical cell membrane; Multi-pass membrane protein. Recycling endosome membrane {ECO:0000250|UniProtKB:O70127}; Multi-pass membrane protein {ECO:0000250|UniProtKB:O70127}. Endosome {ECO:0000250|UniProtKB:O70127}. Cell membrane; Multi-pass membrane protein. Note=Internalized at the canalicular membrane through interaction with the adapter protein complex 2 (AP-2) (PubMed:22262466). At steady state, localizes in the canalicular membrane but is also present in recycling endosomes. ABCB11 constantly and rapidly exchanges between the two sites through tubulo-vesicles carriers that move along microtubules. Microtubule-dependent trafficking of ABCB11 is enhanced by taurocholate and cAMP and regulated by STK11 through a PKA-mediated pathway. Trafficking of newly synthesized ABCB11 through endosomal compartment to the bile canalicular membrane is accelerated by cAMP but not by taurocholate (By similarity). Cell membrane expression is up-regulated by short- and medium-chain fatty acids (PubMed:20398791) {ECO:0000250|UniProtKB:O70127, ECO:0000269|PubMed:20398791, ECO:0000269|PubMed:22262466}

Tissue Location

Expressed predominantly, if not exclusively in the liver, where it was further localized to the canalicular microvilli and to subcanalicular vesicles of the hepatocytes by in situ

Background


Involved in the ATP-dependent secretion of bile salts into the canaliculus of hepatocytes.

References

Strautnieks S.S.,et al.Nat. Genet. 20:233-238(1998).
Mol O.,et al.Submitted (MAR-1999) to the EMBL/GenBank/DDBJ databases.
Hillier L.W.,et al.Nature 434:724-731(2005).
Mochizuki K.,et al.Am. J. Physiol. 292:G818-G828(2007).
Jansen P.L.M.,et al.Gastroenterology 117:1370-1379(1999).

Images

Anti-ABCB11 Antibody (C-term) at 1:2000 dilution + mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 146 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.